



ISSN 2610-931X

CEIS Tor Vergata

RESEARCH PAPER SERIES Vol. 22, Issue 1, No. 573 – April 2024

Caring Connections: Immigrant Caregivers and Long-Term Elderly Care in Italy

Elisa Capretti, Joanna Kopinska, Rama Dasi Mariani and Furio Camillo Rosati

Caring Connections: Immigrant Caregivers and Long-Term Elderly Care in Italy^{*}

Capretti L.¹, Kopinska J. A.^{2,1}, Mariani R. D.^{3,1}, and Rosati F. C.^{4,1} ¹CEIS, University of Rome "Tor Vergata" ²DISSE, Sapienza University of Rome ³Dept. of Law, Roma Tre University ⁴DEF, University of Rome "Tor Vergata"

Abstract

This paper investigates the impact of migrant-provided home-based care on elderly health in Italy, analysing hospitalisation frequency, duration, and mortality. Using an instrumental variable approach to mitigate endogeneity between local health status and migratory flows, we show that migrant-provided home-based care reduces the frequency of hospital admissions (extensive margin) and their duration (intensive margin). Regarding the former, a one percentage point increase in the immigrant-to-elderly population ratio leads to a 4% decrease in long-term and rehabilitation (LR) stays, with no effect on acute stays. Concerning the latter, we find that a similar change in the migrant inflows translates to a 1.5% reduction in admission duration, with LR admissions reaching a 3.3% decline. These effects primarily stem from traumatic injuries, musculoskeletal disease, and genitourinary disorders, particularly linked to home-based mobility and treatment management. Our back-of-the-envelope calculations suggests that a 1.3 percentage point increase in the migrant-to-elderly population ratio registered in our analysis period could potentially reduce LR elderly hospitalisation costs by approximately 8% and yield annual public budget savings equivalent to around 0.59% of total hospitalisation expenditures.

^{*} Funding Declaration: Financial support from the Italian Minister of University, Grant PRIN2017(Prot.2017KHR4) is gratefully acknowledged.

1 Introduction

The impact of immigration on the receiving economy has undergone extensive scrutiny in the academic literature. Most of the attention, however, has been devoted to the changes immigration induces in the labour market. Less attention has been paid to the impact on domestically produced goods and services and on the allocation of non-market time. A large share of low-skilled immigrants specializes in the provision of domestic services that substitute or complement the time devoted by the household members to domestic production.

With the rapid aging of Western societies, the demand for long-term care (LTC) services is increasing. Consequently, many elderly individuals rely on informal care provided by family members, given the limited capacity of the (public) formal care system. Informal caregivers offer assistance with daily tasks, medication management, and other forms of support, which play a vital role in maintaining individuals' health and reducing hospitalisation rates (Barnay & Juin, 2016; Charles & Sevak, 2005; Van Houtven & Norton, 2004).

However, the supply of informal caregiving provided by family members is declining due, inter alia, to the reduction in fertility and the increasing labour force participation, particularly among women. Households are, therefore, increasingly using paid home-based care. In countries like Germany, France, and the UK, paid care is typically provided by NGOs or private firms, while Southern European countries like Spain, Greece, and Italy predominantly rely on domestic services, with a significant portion of the workforce constituted by immigrants. Notably, in Italy, immigrants represent approximately 60 per cent of the total domestic workers,² and 80 per cent of the live-in assistants providing a wide range of household services (Mariani & Rosati, 2022).

The recent inflows of immigrants to Western countries have somewhat addressed the shortfall in publicly provided LTC services. However, their impact on the health outcomes of the elderly has not been studied, nor their effects on the national health system with the exception of Bettin & Sacchi (2020). To address this gap, our study analyses the impact of immigrant-provided home-based LTC services in Italy on various adverse and costly health outcomes, including hospitalisations, their duration, and mortality rates among the elderly population. We also offer rough estimates of the resulting potential savings for the public health system that informal care provision could generate.

The connection between home-based care and hospitalisation is intricate, influenced by factors such as the health condition of the care recipient, the availability and competence of non-specialized home caregivers, and the quality and availability of alternative care options. Non-specialized homebased care emerges as a pivotal element in identifying and managing a broad spectrum of chronic conditions among the elderly, as demonstrated by Kemper et al. (2008). In particular, home-based

² <u>https://data.oecd.org</u> (last accessed on April 5th, 2024)

care plays a critical role in early symptom detection, thereby allowing timely management of the disease. Furthermore, it promotes mobility, encourages physical activity and healthy eating habits, supports medication compliance, and enables vigilant monitoring of symptoms. Additionally, home-based care is instrumental in facilitating individuals' transition from hospital settings back to their homes following a hospitalisation episode (Costa-Font et al., 2018).

Beyond these direct health benefits, home-based care plays a crucial role in preventing unmet needs within formal healthcare. Notably, Rice et al. (2009) show that home-based care can mitigate the inefficient utilisation of long-term hospital services. Similarly, Weaver & Weaver (2014) find that the presence of informal care, while not necessarily reducing the likelihood of hospitalisation, significantly shortens thelength of stays (LoS) in hospital.

For our analysis, we leverage the Hospital Discharge Data ("*Schede di Dimissione Ospedaliera*" – SDO) which encompasses all hospitalisations across public and private facilities in Italy from 2006 to 2015 at the province level (NUTS-3). The SDO provides detailed information on both the sociodemographic and clinical aspects of each hospitalisation, such as diagnosis, procedures, duration, and tariffs that allow the calculation of the relevant costs. We combine the health data with the distribution of immigrants of age between 15 and 64 as a ratio of the native population over 65 across Italian provinces over the same period.

To establish the causal link between immigration and the health outcomes of the elderly, one must inevitably deal with the very likely non-random allocation of immigrants across provinces. Newlyarrived immigrants might select to areas where demand for the services they supply is high. We thus employ an instrumental variable (IV) estimation, based on a version of the shift-share instrument à la Card (first introduced by Altonji & Card, 1991 and Card, 2001). To do so, we ensure that our IV, which reflects the past distribution of immigrants across Italian provinces, withstands rigorous validity tests recommended by recent literature on Bartik instruments (Borusyak et al., 2022; Goldsmith-Pinkham et al., 2020) and confirm the plausibility of our instrument. To further test the soundness of our research design, we exploit the long-difference variation of the distribution of immigrants and of the outcome variables over the period of our analysis (between 2006 and 2015). Then, to discard reverse causality, we also show that, in a placebo fashion, the instrument-predicted change of immigrants over the period does not correlate with the variation in our main outcome variables in a previous period (between 2001 and 2005). Lastly, we confirm the robustness of our results controlling for lagged changes in outcome variables (between 2001 and 2005) as in Dix-Carneiro et al. (2018). The latter specification is another way to address concerns about pre-existing trends in hospitalisations and average LoS that could be correlated with (future) migration shocks.

We show that home-based care provided by immigrants has the potential to generate savings in formal healthcare expenditure through two distinct channels. Firstly, impacting the extensive margin, we show that an increase of one percentage point in the ratio between the number of working-age immigrants and the native population over 65 leads to a reduction of around 118 LR hospitalisations per 100 thousand elderly residents, representing roughly 4% of the baseline average. We find no statistically significant effects on the frequency of acute hospitalisations. Secondly, in addition to the reduced hospitalisation frequency, migrant home-based care also leads to shorter average hospital stays, influencing the intensive margin. We show that a one percentage point increase in the availability of migrant-provided home-based care translates to a 1.5% decrease in the mean duration of all hospital admissions. This impact is notably amplified in the case of long-term and rehabilitation hospital stays, reaching a substantial reduction of 3.3%. When scrutinising the effect by diagnostic groups, we show that our effect is driven by hospitalisations originating in traumatic injuries, musculoskeletal disease, and genitourinary disorders.

Considering both the intensive and extensive margins, our rough estimation, conducted through a back-of-the-envelope calculation, indicates that the annual increase in the migrant-to-elderly population ratio (averaging 1.3 percentage points annually during our analysis period) accounts for roughly 8% of the long-term and rehabilitation hospitalisations dedicated to the elderly observed in the study. Extrapolating from this, it is plausible to infer that the influx of immigrants over our study period may have resulted in annual savings to the public budget equivalent to approximately 0.59% of total expenditures on hospitalisation. Looking towards the long-term perspective, particularly considering Italy's significant pace of population aging, these potential savings are likely to become increasingly important and could rise by a third.

2 Long-term care for the elderly in Italy

According to the OECD, Italy's health expenditure in 2019, covering all functions, amounted to 8.7% of GDP, in line with both the OECD and the EU averages. In the same year, Italy allocated approximately 1.7% of its GDP to public spending on LTC, slightly exceeding the EU average of 1.6%. While these figures remained stable in the years leading up to 2019, they showed a significant increase in 2020 due to the Covid-19 pandemic,³ posing a great challenge to the sustainability of the public LTC.

Italian public LTC is structured around two schemes: public provision of services and cash transfers. Cash transfers account for a large share of the funding (52% in 2019) and consist of monthly

³ https://www.oecd.org/health/

payments disbursed to individuals with disabilities or severe dependency, irrespective of their income (NNA, 2021).⁴ Financed through general taxation and managed by the central government, they provide assistance to roughly 11% of the population aged 65 and above.

Public provision of services is managed by local health authorities and encompass Integrated Home Care (Assistenza Domiciliare Integrata - ADI), and Home Care Service (Servizio di Assistenza Domiciliare - SAD). ADI provides medical, nursing, and rehabilitation services to individuals lacking self-sufficiency, facing frailty, and/or suffering from chronic illnesses. On the other hand, SAD is primarily dedicated to the essential needs of the elderly population, such as personal hygiene, dressing, mobility, and medication management. The coverage rate of these services is significantly lower compared to cash transfers, with residential care reaching approximately 3% of the potential beneficiaries and home-based care around 5% of the target population. The challenge of ensuring sufficient public home-based care for the elderly is particularly pronounced in Southern regions, where the scarcity of formal public services and low employment rates compel families to take on the responsibility of organising alternative care arrangements, increasingly relying on informal caregivers (Melchiorre et al., 2021).

In Italy women represent about 57% of informal LTC providers, frequently reducing their working hours or exiting the workforce to care for their relatives (Peri et al., 2015). In 2020, informal caregivers accounted for around 10% of Italy's total population and 13% of its labour force, devoting roughly 20 hours per week to caregiving duties (NNA, 2021).

Although the public sector offers support to informal caregivers, such as paid leave, tax deductions, and vouchers for hiring domestic workers, these measures are often inadequate and unequally distributed across regions. Consequently, many informal caregivers rely on paid domestic work to supplement or replace the time they dedicate to elderly care. As of 2015, the last year of our sample, data from the Italian National Institute for Social Security (Istituto Nazionale Previdenza Sociale – INPS), analysed by DOMINA, revealed the presence of 905,224 regular domestic workers, of which more than 75% were immigrants and approximately 85% were women (DOMINA, 2022). The main regions of origin of the foreign domestic workers are Eastern Europe, accounting for approximately 45% of the sectoral workforce, and Asia, accounting instead for more than 15% (see Figure 1). Furthermore, irregular workers add approximately 1 million to the domestic workforce, constituting 52.3% of the sector's total workforce, significantly surpassing the overall average irregularity rate of 12% (DOMINA, 2022).

⁴ Cash transfers for LTC are not subject to conditionalities on the use but only on the assessed need.




Figure 1: Domestic Workers in Italy (2006-2015)

Notes: Author's elaboration on INPS data obtained from "DOMINA National Observatory on Domestic Work " (DOMINA, 2022). The definitions domestic workers follow the ILO standard classifications, referring to individuals who perform work in or for a private household or households on a full-time or part-time basis as live-in or live-out workers.

The two peaks in the series of regular domestic workers, both total and immigrants, shown in Figure 1, occur in 2009 and 2012 and coincide with two regularisation laws for irregular immigrant domestic workers already residing in the country.

Recent studies focusing on the immigrant populations in the United States and Germany have observed an increasing specialisation of immigrants in delivering institutional care within nursing homes. Furtado & Ortega (2023) and Grabowski et al. (2023) document that a higher presence of immigrant among nursing home workers results in increased utilisation rates and better health outcomes for residents, particularly in settings with a higher proportion of Hispanic staff. However,

to the best of our knowledge, the role of immigrants in providing paid home-based care has received no attention so far.

In Italy, immigrants have significantly affected the provision of domestic informal care services, thus affecting the demand for institutionalised LTC and complementing the efforts of household members involved in caregiving and domestic activities. We will make use of the variation in the availability of domestic services across space and time, due to differences in the number of working-age immigrants in Italy, to study the effect of paid home-based services on elders' health.

3 Data

The core of our analysis is conducted using Hospital Discharge Data (SDO) spanning from 2004 to 2015, collected by the Italian Ministry of Health. The data refer to all hospitalisations within public and publicly-funded private hospitals. Italy's universal public healthcare system offers free hospitalisations, eliminating cost barriers and selection mechanisms, providing an ideal setting for the analysis.

The SDO data provide extensive information about hospitalisations, including their type, the primary and secondary diagnosis, procedures, and discharges, as well as various socio-demographic characteristics such as age, gender, nationality, place of birth, and residence.

We focus on the hospitalisations of Italian citizens aged 65 and over, among which we distinguish between Acute and Long-term/Rehabilitation (LR) admissions. Acute hospitalisations address immediate and critical medical needs and are typically of short duration. Conversely, LR hospitalisations provide prolonged and comprehensive care, often spanning long treatment periods addressing chronic illnesses or complex medical conditions.

We focus on both the incidence and the duration (Length of Stay - LoS) of Acute and LR hospitalisations. In particular, we consider the ratio of both kinds of hospitalisations to the native population aged 65 and above for each province-year pair. The average LoS is computed by dividing the total number of hospitalisation days in a province-year pair by the number of hospitalisation events.

Based on the International Statistical Classification of Diseases and Related Health Problems (ICD-9) codes, we group the hospitalisations in eight major diagnostic groups.⁵ For the disease-specific subgroups of hospitalisations, we compute the frequency and the average LoS as discussed above. Finally, as an additional and comprehensive health outcome, we rely on age-standardized mortality data obtained from the database Health for All by the Italian National Statistical Office (Istat). Our

⁵Mental Disorders, Cancers, Nervous System Disorders, Circulatory Disorders, Respiratory Disorders, Genitourinary Disorders, Musculoskeletal Disorders, and Traumatic Injuries.

analysis focuses on mortality among individuals aged 75 and older, available for the years 2003-2016.

As shown in Table 1 (Panel a), the annual average of hospital discharges per 1,000 residents aged 65 and older is 395. Acute cases show an average of 367 cases, while LR discharges constitute a significantly smaller proportion averaging nearly 28 stays per 1,000 residents above 65 years of age. On average, hospitalisations last around 10 days, with acute stays lasting about 9 days and long-term/rehabilitation admissions averaging 29 days. According to Panel (b) of Table 1, age-standardized mortality among individuals aged 75 and older averages 643 deaths per 10,000 residents for women and 790 deaths for men, with the overall mortality rate being 716 deaths per 10,000 residents.

Variable	Obs.	Mean	St. Dev.
Panel a: Hospitalisations			
Hospitalisations per capita (All cases)	1030	0.395	0.073
Hospitalisations per capita (Acute)	1030	0.367	0.0709
Hospitalisations per capita (LR)	1030	0.028	0.013
Average Length of stay (All cases)	1030	10.299	1.441
Average Length of stay (Acute)	1030	8.635	0.926
Average Length of stay (LR)	1030	29.005	6.285
Panel b: Mortality			
Mortality (total)	1030	716.301	42.503
Mortality (women)	1030	643.344	41.981
Mortality (men)	1030	789.494	49.554

Table 1: Descriptive statistics on hospitalisations and mortality

Notes: The data refer to 103 provinces spanning the time period from 2006 to 2015. Statistics in Panel (a) are derived from an initial sample of hospital discharges within the entire population of Italian citizens aged 65 and older, accounting for approximately 4 million hospitalisations annually. Statistics are weighted by the relevant province 65+ population size. Data in Panel (b) refer to official ISTAT province-level age-standardized mortality for 10,000 individuals aged 75 and older for the same period.

The main explanatory variable included in the analysis is the number of immigrants of age between 15 and 64 as a ratio of the native population over 65 obtained from the administrative data collected by the Italian statistical institute (Istat). Additionally, we employ a set of province-level controls that account for the time-varying local socio-economic and demographic characteristics: per-capita

income, the share of widows and widowers in the population over 65.⁶ Table 2 presents the relative summary statistics.

We combine hospitalisation and mortality data with the administrative data on the number of immigrants and other characteristics, creating a balanced panel of 103 provinces over the period 2006-2015, for a total of 1030 observations.

Variable	Obs.	Mean	St. Dev.
Share of immigrants	1030	24.2	0.118
Instrumented Share of Immigrants	1030	33.8	0.300
Income pc (log)	1030	13.726	0.830
Widowers (%)	1030	1.633	0.725
Widows (%)	1030	2.287	0.957
Population over 65 years (weight variable)	1030	113.741	115.081

Table 2: Descriptive statistics of migration and other province-level data

Notes: The statistics refer to 103 provinces spanning the time period from 2006 to 2015. The number of immigrants is expressed as a ratio to the resident population aged 65 and older and reported in the table for 100 individuals. The statistics concerning immigration are weighted by province 65+ population size. Population 65 and older is used as weight variable, and it is expressed in thousands.

4 Empirical analysis

4.1 Conceptual Framework

Framing our analysis in the Grossman (1972) model, we assume that investment in health is obtained through the use of three inputs: formal care, informal care (provided by relatives), and non-specialized home-based care (provided through services bought on the market). For the latter, we refer in particular to the services provided by hired non-specialized personnel (co-resident or not).

In the case of informal care provided by relatives, the price is given by the opportunity cost of time, while the price of formal care might include, besides the out-of-pocket expenditures, the monetary value of the disutility of being in an institution rather than at home.

⁶ In a further robustness test we also control for the share of native women aged between 45 and 64 as ratio of native population over 65. According to NNA (2022) this group consists of about 57% of total informal caregivers in Italy. Therefore, with this specification, we show that the impact of home-based care is robust also when we net out the presence and availability of informal caregivers.

If we assume a unitary household model, and if the inputs are all substitutes, a reduction in the price of home-based care provided by hired workers will reduce the demand for the other inputs, including formal care. It will also increase the overall investment in health, if the latter is not an inferior good.

Clearly, one could think of more complex production structures that might also give rise to interactions across inputs considering nested production functions. For example, we might consider informal and home-based care as complementary inputs in the health investment function, with informal and home-based care produced with the time of relatives and hired helpers, respectively. However, in the absence of data on the time devoted by relatives to informal care, we cannot investigate this further. In the estimates, however, we use controls for taking into consideration the characteristics of the average family composition.⁷

Because of the data constraint just mentioned, we do not consider the possibility of strategic interactions between the household members in the provision of informal care, which has been discussed in some papers (*e.g.* Byrne et al. (2009)). In the model developed by Byrne et al. (2009), for example, the spouse is more likely to provide care with respect to adult children because of the lower burden experienced by spouses in the caregiving role.

In what follows, we thus carry out reduced form estimates of the impact of home-based care provided by hired migrant workers on the use of formal care, as proxied by the hospitalisation episodes and duration, and on the health status of the elders, proxied by the age-standardized mortality rate.

4.2 Identification strategy

The presence of immigrants is likely to be endogenous to the use of formal care and the health status of the elderly population. Immigrants might self-select based on their willingness and ability to provide quality healthcare in locations with growing demand for domestic workers, associated with the increasing needs of the elderly. This is likely to exert an attenuation bias on the estimation of the impact of the arrival of immigrants on the health status of the elders.

A similar and mirrored selection mechanism could drive the elders in need of assistance to move to places where the supply of non-institutionalized care is growing. Conversely, immigrants and the elderly might both be influenced by changes in local conditions such as socio-economic factors, neighbourhood characteristics, or healthcare infrastructures. For example, provinces with worsening economic prospects might attract fewer immigrants seeking employment and, at the same time, feature less healthy elderly individuals.

⁷ In particular, we consider the initial shares of widows and of widowers as proxies for (the lack of) family members that might complement the home-based care provided by immigrant workers.

While province and year-fixed effects absorb time-invariant effects and common time trends in the estimates, the issue of a possible bias arising from time-varying omitted variables affecting both immigrant flows and health conditions concurrently remains a concern.

The mobility of individuals over 65 is very limited in Italy, with a mere 0.8 percent changing their main residence in 2017⁸. Consequently, we can consider their location to be exogenous with respect to the availability of informal care. However, this assumption does not hold for immigrants, who select their location upon arrival and exhibit a relatively high rate of subsequent mobility (Mariani et al., 2023).

To mitigate potential endogeneity stemming from immigrants' location and time-varying omittedvariables, we adopt an instrumental variable (IV) approach. Specifically, we use a version of the Card instrument, pioneered by Altonji and Card (1991) and further developed by Card (2001), which exploits the different timing and size of migrant inflows by origin, distributing them across provinces on the basis of the 1991 shares of immigrants by origin. As shown in recent literature (Borusyak et al., 2021; Goldsmith-Pinkham et al., 2020) this *shift-share* instrument can be considered exogenous, conditionally on a set of validity tests that we perform and discuss in section 5.3.

More precisely, our instrument uses the shares observed in 1991 of immigrants coming from ten different world regions (North Africa, Other Africa, East Europe, Asia, South America, Oceania, North America, and Western Europe) and is defined as:

$$Z_{i,t} = \sum_{m} \lambda_{i,1991}^{m} \times Imm_{ITA,t}^{m} \tag{1}$$

where $\lambda_{i,1991}^m$ is the share of immigrants (males and females) observed in 1991 in province *i* from region of origin *m*, and $Imm_{ITA,t}^m$ is the stock of working-age immigrants in Italy at time *t* from region of origin *m*.

Consequently, the second-stage equation takes the following form:

$$H_{i,t} = \beta_1 \widehat{M_{i,t}} + \beta_2 X_{i,t} + \lambda_i + \tau_t + \epsilon_{i,t}$$
⁽²⁾

where $H_{i,t}$ denotes one of the health outcomes (hospitalisations frequency, average LoS, and mortality) in province *i* and year *t*. $\widehat{M_{i,t}}$ indicates the instrumented share of immigrants aged 15-64 as a ratio of the population aged 65 and older, and $X_{i,t}$ is a vector of control variables at the province level, including per-capita income, the percentage of widows, and the percentage of widowers. λ_i and τ_t are province and time fixed effects, respectively.

⁸ <u>https://demo.istat.it/</u> (last access April, 8 2024)

We test the validity of our instrumental-variable approach in several ways, many of which leverage the variation in both immigrant share and health outcomes over the long difference period, namely the overall change between 2006 and 2015. Therefore, we first show that the long-difference specification between 2006 and 2015 yields comparable results to the main specification, which instead exploits yearly variations.

Using this result as a benchmark, we mitigate reverse causality concerns by controlling for the fact that the instrument-predicted change in immigrant share during our analysis period (2006-2015) does not explain variations in the main outcome variables in the pre-sample period. Additionally, we explicitly test for the exogeneity of the initial (1991) immigrant shares, which may potentially be endogenous to local economic and health conditions. In particular, it is crucial to rule out any spurious correlation between past shares and persistent trends in local economic and health conditions. To this aim, following Goldsmith-Pinkham et al. (2020), we check whether the 1991 shares are exogenous and not correlated with pre-treatment trends.

Lastly, we confirm the robustness of our results by controlling for lagged changes in outcome variables (between 2001 and 2005 for hospitalisations frequency and average LoS and between 2003 and 2005 for mortality), as in Dix-Carneiro et al. (2018). This specification helps mitigate concerns regarding pre-existing trends in the health status of the elderly that could be correlated with future migration shocks.

5 Results

5.1 Instrumental variable results

As *prima facie* evidence of the relevance of the instrument, we show the output of the first-stage regression. As displayed in Table 3, the coefficient of the instrumental variable described in equation (1) in section 4.2, is positive and significant, and the weak IV test (the Kleibergen- Paap rk Wald F statistics) is high at around 39.

	Share of immigrants
Instrumented share of immigrants	0.184***
-	(0.030)
Observations	1030
Number of provinces	103
Year FE	YES
Province FE	YES
Controls	YES

Table 3: First-Stage Regression

Notes: Authors' own calculations on Health Ministry's and ISTAT data. The reported coefficient comes from a panel fixed effects model at the province level for the years 2006-2015. The dependent variable is the ratio between the working-age immigrant population over the native population aged 65 and older. The control variables include: the (log of) income per capita, and female and male widow shares (%). The regression is weighted by the province population 65+. Robust standard errors in parenthesis with p < 0.1 * p < 0.05 * p < 0.01

The exclusion restriction of the instrument hinges on the assumption that the 1991 distribution of immigrants by area of origin across provinces is not correlated with hospitalisation and mortality in the period under analysis (2006-2015), except through its impact on current immigration. As mentioned earlier, we carry out several tests to substantiate this assumption, detailed in section 5.3. To begin, we present the results of the 2SLS estimations assessing the impact of immigrant workers on the health conditions of the elderly⁹.

5.1.1 Hospitalisations

The coefficient estimates reported in Table 4 suggest that the presence of migrants does not affect the overall hospitalisation rate, however, when we disaggregate them by Acute and Long-term hospitalisations, we find that the latter are significantly and negatively affected by the presence of immigrants. An increase of one percentage point in the ratio between the number of working-age immigrants and the native population over 65 (e.g., from 0.242 to 0.252) leads to a reduction of around 118^{10} LR hospitalisations per 100 thousand elderly residents, representing roughly 4% of

⁹ In appendix we show that the results are confirmed also when adding a control variable for infomal care provided by family members.

¹⁰ This is computed by multiplying the coefficient by 0.01 and then by 100,000 residents.

the baseline average. This is not surprising as Acute hospitalisations typically address conditions that require specialized medical care and can be hardly substituted by non-specialised home-based care. On the contrary, for Long-term care and, to a certain extent, rehabilitation, home-based care can represent a viable substitute. In fact, the elderly often resort to hospitalisation when they lack basic support to deal with their ailments at home.

Importantly, the OLS estimates presented in Table A1.1 of the appendix indicate the presence of an upward bias. Hence, without addressing the endogeneity between local elderly health and immigration inflows, the OLS results suggest that immigrants tend to settle in provinces where acute hospitalisations among the elderly are more frequent, while having no significant relationship with long-term hospitalisation episodes. This pattern suggests a selection mechanism, wherein migrants tend to settle in provinces where the health status of the elderly is relatively poorer and the demand for their services higher.

	(1)	(2)	(3)
	Acute & LR	Acute	LR
Share of immigrants	0.0603	0.178	-0.118***
	(0.229)	(0.227)	(0.024)
Income per capita (log)	0.115	0.107	0.009
	(0.0965)	(0.088)	(0.010)
Widowers (%)	0.0664**	0.064**	0.003
	(0.0299)	(0.029)	(0.003)
Widows (%)	-0.0154	-0.016	0.001
	(0.0240)	(0.023)	(0.002)
Observations	1,030	1,030	1,030
R-squared	0.675	0.685	0.121
Number of provinces	103	103	103
Year FE	YES	YES	YES
Province FE	YES	YES	YES
F Stat	38.604	38.604	38.604

Table 4: IV results for the number of hospitalisations broken down by type

Notes: Estimates of equation 2. For each hospitalisation type, the dependent variable is calculated as the number of relevant hospitalisations over the native population 65+. In column 1 we include all hospitalisations, in column 2 we focus on acute hospitalisations, and in column 3 on long-term and rehabilitation. The endogenous regressor is the ratio between the working-age immigrant resident population over the size of native population aged 65 and older. Each regression is weighted by the province population 65+. The reported F Statistics is the Kleinbergen–Paap rk Wald F statistics. Robust standard errors in parenthesis with p < 0.1 * p < 0.05 *** p < 0.01.

5.1.2 Average Length of Stay (LoS)

Interestingly, the average LoS in the hospital setting is reduced by the presence of immigrants, as shown in Table 5. An increase of one percentage point (e.g., from 0.242 to 0.252) in the ratio between the working-age immigrants and the native population over 65 leads to a reduction of around 0.14 days in hospital stays, which represents about 1.5% at the mean duration (with the average being 10 days). The effect holds for both kinds of hospitalisations but is particularly pronounced for long-term care. The reduction amounts to 0.96 for LR hospitalisations, roughly 3.3% relative to the average duration of around 29 days. For Acute cases, the reduction is 0.03 days, or 0.3% of the average duration of 9 days.

Indeed, the duration of LR hospitalisation episodes appears to be most responsive to the availability of home-based care. This is because hospital discharges can happen sooner if elderly individuals receive assistance at home for medical treatments, basic daily activities, and outpatient medical visits, thereby reducing the necessity for prolonged hospital stays.

When examining the different causes of hospitalisations mentioned earlier (as detailed in Section 5.2.2), it is evident that the observed effect is primarily driven by a subset of causes. For instance, there is a significant reduction of around 11 days in the length of stay for hospitalisations with primary diagnoses related to traumatic injuries, respiratory disease, and genitourinary disorders.

A reduction in the length of stay in the hospital was also found in Weaver and Weaver (2014) for informal care provided by family members. Our results strengthen and extend this evidence for the case in which care is provided by immigrants as a substitute for family members.

	(1)	(2)	(3)
	Acute &		
	LR	Acute	LR
Share of immigrants	-14.17***	-3.033**	-95.66***
	(2.202)	(1.240)	(19.74)
Income per capita(log)	-1.483**	-0.668	-8.784*
	(0.633)	(0.459)	(4.615)
Widowers (%)	-0.806**	-0.735***	-1.565
	(0.321)	(0.250)	(2.072)
Widows (%)	0.0486	-0.0121	-0.236
	(0.261)	(0.248)	(1.466)
Observations	1,030	1,030	1,030
R-squared	0.076	0.153	-0.037
Number of provinces	103	103	103
Year FE	YES	YES	YES
Province FE	YES	YES	YES
F Stat	38.604	38.604	38.604

Table 5: IV results for the LoS of hospitalisation broken down by type

Notes: Estimates of Equation 2. For each hospitalisation type, the dependent variable is calculated as the total hospitalisation days over the number of relevant hospitalisations. In column 1, we refer to all hospitalisations, in column 2, to acute hospitalisations, while in column 3, to long-term and rehabilitation hospitalisations. The endogenous regressor is the ratio between the working-age immigrant resident population over the size of native population aged 65 and older. Each regression is weighted by the province population 65+. The reported F Statistics is the Kleinbergen–Paap rk Wald F statistics. Robust standard errors in parenthesis with * p<0.1 ** p<0.05 *** p<0.01.

5.1.3 Mortality

When examining the impact of home-based care availability on the health of the elderly, we also analyse mortality rates. The findings presented in Table 6 suggest that the presence of immigrants is likely to reduce the mortality rate, with the results mainly driven by the effect on women. The estimates show relatively weak statistical significance, yet indicate a consistent direction of health improvements associated with the availability of immigrant workers. Beyond managing medical and clinical treatments, caregivers are also inclined to encourage healthy lifestyles. Moreover, they provide emotional support, thereby fostering overall well-being. The gender gradient observed in the results is likely attributed to the more favourable survival pattern of women compared to men among the

population aged 75 and over. On average, women tend to have a higher life expectancy, making them more likely to benefit from home-based care at older ages.

	(1)	(2)	(3)
	Total	Female	Male
Share of immigrants	-189.0*	-235.4**	-142.7
	(106.4)	(110.8)	(116.5)
Income per capita(log)	-6.433	-16.52	3.653
	(34.77)	(34.35)	(41.02)
Widowers (%)	-9.409	1.852	-20.67
	(15.69)	(12.41)	(21.09)
Widows (%)	-9.999	-10.78	-9.216
	(11.46)	(9.196)	(15.12)
Observations	1,030	1,030	1,030
R-squared	0.438	0.454	0.428
Number of provinces	103	103	103
Year FE	YES	YES	YES
Province FE	YES	YES	YES
F Stat	38.604	38.604	38.604

Table 6: IV results for mortality rates

Notes: Estimates of Equation 2. The dependent variable is represented by agestandardized mortality for 10,000 population aged 75 and older. The endogenous regressor is the ratio between the working-age immigrant population over the size of native population aged 65 and older. The reported F Statistics is the Kleinbergen– Paap rk Wald F statistics. Each regression is weighted by the province population 65+. Robust standard errors in parenthesis with *p<0.1 **p<0.05 ***p<0.01.

5.2 Results by Diagnostic Groups


To provide a more comprehensive analysis and validate our findings, we disaggregate the long-term care hospitalisations into eight major Diagnostic Groups.

5.2.1 Hospitalisations

Figure 2 and Table 7 show that an increase in the share of immigrants leads to a reduction of LR hospital stays for diagnoses related to mental issues, nervous system disorders, genitourinary disorders, musculoskeletal disorders and traumatic injuries. Specifically, we can observe a reduction

of 51 cases every 100,000 residents related to genitourinary system disorders (column 6 in Table 8) as having the assistance of caregivers can indeed help elders with better hygiene practices. Additionally, as indicated by Rogers et al. (2008) and Lean et al. (2019), urinary infections, which are common among the elderly, may occur due to factors such as limited mobility and dehydration, both of which are likely to be alleviated in the presence of home-based care. The promotion of mobility is further evidenced by the observed reduction in hospitalisations attributed to musculoskeletal disorders and traumatic injuries, estimated at approximately 32 and 19 cases per 100,000 residents, respectively. To a more limited extent, the presence of immigrant caregivers can also reduce hospitalisations due to mental disorders, by 3 cases every 100,000 residents. This reduction likely reflects the positive impact of home-based care to emotional well-being and medication compliance, reducing the risk of depression, as suggested by Barnay and Juin (2016) in the context of informal care. Conversely, we find no effect on the frequency of hospitalisations due to the more severe and acute nature of such hospitalisation events, potentially less influenced by home-based care.





Notes: The coefficient estimates come from eight distinct IV regressions at the province-year level, each for one dependent variable defining long-term hospitalisation episodes for 8 major diagnostic groups: mental, cancers, nervous system, respiratory, circulatory, genitourinary, musculoskeletal, and traumatic injuries. The explanatory variable is the ratio between the working-age immigrants over the native population aged 65 and older. Each regression controls for province and year fixed effects and is weighted by the province population 65+. The reported F Statistics is the Kleinbergen–Paap rk Wald F statistics. Robust standard errors in parenthesis p < 0.1 * p < 0.05 * p < 0.01

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
	Mental	Cancers	Nervous	Respiratory	Circulatory	Genitourin.	Musculoskel.	Traumatic
Share of immigrants	-0.00316*	0.000841	-0.0150***	-0.00165	-0.00486	-0.0510***	-0.0326***	-0.0192**
Share of miningrams	(0.00170)	(0.00149)	(0.00378)	(0.00357)	(0.00651)	(0.00791)	(0.00626)	(0.00854)
Income per capita (log)	-0.00106*	0.000868**	-0.00116	0.000286	0.00140	0.00376	0.00207	0.00124
1 1 0	(0.000560)	(0.000432)	(0.00124)	(0.000575)	(0.00191)	(0.00320)	(0.00179)	(0.00174)
Widowers (%)	-5.84e-05	-0.000108	-0.000398	0.000805	0.00116	0.000207	-0.000602	0.000494
	(0.000239)	(0.000149)	(0.000390)	(0.000514)	(0.000965)	(0.00106)	(0.00105)	(0.000471)
Widows (%)	0.000519**	9.67e-05	0.000266	-0.000447	-0.000963	0.00102	0.000975	0.000339
	(0.000258)	(9.44e-05)	(0.000284)	(0.000380)	(0.000695)	(0.000791)	(0.000818)	(0.000429)
Observations	1,030	1,030	1,030	1,030	1,030	1,030	1,030	1,030
R-squared	0.009	0.120	0.065	0.108	0.169	0.064	0.174	0.029
Number of provinces	103	103	103	103	103	103	103	103
Year FE	YES	YES	YES	YES	YES	YES	YES	YES
Province FE	YES	YES	YES	YES	YES	YES	YES	YES
F Stat	38.604	38.604	38.604	38.604	38.604	38.604	38.604	38.604

Table 7: IV results for LR hospitalisations by major diagnostic groups


Notes: The dependent variables are long-term cases for 8 major diagnostic groups: mental (column 1), cancers (column 2), nervous system (column 3), respiratory (column 4), circulatory (column 5), genitourinary (column 6), musculoskeletal (column 7) and traumatic (column 8). The explanatory variable is the ratio between the working age immigrants over the native population aged 65 and older. Each regression is weighted by the province population 65+. The reported F Statistics is the Kleinbergen–Paap rk Wald F statistics. Robust standard errors in parenthesis p < 0.1 * p < 0.05 * p < 0.01

19

5.2.2 Average length of stay (LoS)

Concerning the length of stay of LR stays, Figure 3 and Table 8 show a significant reduction in the duration of stays due to nervous system disorders, respiratory system disorders, circulatory disorders, genitourinary system disorders, musculoskeletal system disorders, and traumatic injuries. In particular, the presence of immigrant home-based care is found to reduce the duration of hospital stays for traumas by around 1 day¹¹, which is likely related to home-based rehabilitation, exercise, and mobility. The duration of hospital stays appears to be also reduced for disorders related to the respiratory system (by 1.2 days) and cardiovascular issues (by 0.54 days). While the two diagnostic groups do not appear to be affected by the provision of home-based care in terms of the likelihood of hospitalisation, regarding duration, we find sizeable effects likely due to the substitutability between the hospital setting and home-based care in case of issues such as medication management, oxygen therapy, monitoring, and clinical control visits.





Notes: The coefficient estimates come from eight distinct IV regressions at the province-year level, each for one dependent variable defining the length of long-term hospitalisation episodes for 8 major diagnostic groups: mental, cancers, nervous system, respiratory, circulatory, genitourinary, musculoskeletal, and traumatic injuries. The explanatory variable is the ratio between the working-age immigrants over the native population aged 65 and older. Each regression controls for province and year fixed effects and is weighted by the province population 65+. The reported F Statistics is the Kleinbergen–Paap rk Wald F statistics. Robust standard errors in parenthesis p < 0.1 * p < 0.05 *** p < 0.01

¹¹ This number is computed multiplying the coefficient in Table 8 (column 8) by 0.01 (1 percentage point increase in immigrant share).

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
	Mental	Cancers	Nervous	Respiratory	Circulatory	Genitourin.	Musculoskel.	Traumatic
Share of immigrants	120.5	-64.11	-47.87**	-120.2***	-53.98***	-91.83***	-59.43**	-106.0***
	(157.1)	(45.22)	(21.51)	(32.47)	(17.68)	(19.14)	(27.13)	(19.98)
Income pc (log)	-18.80	-1.341	-3.125	-8.502	-6.153	-10.01**	-4.805	-6.266
	(21.83)	(13.95)	(7.230)	(5.169)	(5.257)	(4.057)	(4.188)	(4.927)
Widowers (%)	30.45*	-1.373	-6.209	-0.352	-1.363	-3.625	-0.900	-4.020
	(15.99)	(4.432)	(4.265)	(2.244)	(2.819)	(2.672)	(2.298)	(3.558)
Widows (%)	-18.50	-2.048	1.491	0.309	0.361	0.316	-0.477	1.142
	(11.91)	(2.468)	(3.071)	(1.579)	(1.822)	(1.963)	(1.630)	(2.732)
Observations	1,030	1,030	1,030	1,030	1,030	1,030	1,030	1,030
R-squared	-0.003	0.051	0.026	-0.093	-0.004	0.013	0.056	0.015
Number of provinces	103	103	103	103	103	103	103	103
Year FE	YES	YES	YES	YES	YES	YES	YES	YES
Province FE	YES	YES	YES	YES	YES	YES	YES	YES
F Stat	38.604	38.604	38.604	38.604	38.604	38.604	38.604	38.604

Table 8: IV results for LR average LoS by diagnostic groups

Notes: The dependent variables are LR average LoS for 8 major diagnostic groups: mental (column 1), cancers (column 2), nervous system (column 3), respiratory (column 4), circulatory (column 5), genitourinary (column 6), musculoskeletal (column 7) and traumatic (column 8). The explanatory variable is the ratio between the working age immigrants over the native population aged 65 and older. Each regression is weighted by the province population 65+. The reported F Statistics is the Kleinbergen–Paap rk Wald F statistics. Robust standard errors in parenthesis * p < 0.1 ** p < 0.05 *** p < 0.01

21

5.3 Robustness tests for panel IV results

To conduct validity tests of our instrumental variable and assess the robustness of our estimates to alternative specifications, we require data preceding our analysis period. Specifically, for the period 2001 - 2003, such data are only available at the regional level. To bridge this gap, we employ a multivariate imputation approach, leveraging regional data and other controls used in the main regressions to impute provincial-level data for those years. Regarding mortality data, ISTAT provides standardised mortality rates at the province level from 2003 to 2015, enabling us to run similar validity tests considering 2003 to 2005 as the pre-period.

5.3.1 Validity of the shift-share instrument

The shift-share instrument has been widely used in analysing migration-related economic issues. The exclusion restriction, while not directly testable, depends on the assumption that the distribution of 1991 is not correlated with the outcomes of interest during the analysis period (2006-2015), except through its impact on current migration, known as the *network effect*.

Firstly, we exploit the variation in immigrant shares in the long difference between 2006 and 2015 to ensure comparability with the set of validity tests we aim to perform. We regress the 2006-2015 change in the main outcomes (LR hospitalisations, acute hospitalisations, average LoS, and mortality rate) on the change in predicted immigrant shares over the same period, employing both OLS and 2SLS methods. Table 9 shows that results are similar to those obtained using yearly variation.

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
	Change in	LR cases	Change in	Acute cases	Change	in Los (all)	Change in t	ot. mortality
	OLS	IV	OLS	IV	OLS	IV	OLS	IV
Change in the share of immigrants	0.105***	0.045	0.002	0.393	-5.400*	-11.998**	-216.189**	-314.965**
	(0.026)	(0.053)	(0.106)	(0.296)	(2.977)	(4.704)	(102.040)	(149.402)
Observations	103	103	103	103	103	103	103	103
R-squared	0.200	0.157	0.076	-0.059	0.215	0.134	0.197	0.187
Controls	YES	YES	YES	YES	YES	YES	YES	YES
F Stat	-	19.89	-	19.89	-	19.89	-	19.89

Table 9: Long-Difference Results (2006-2015)

Notes: The estimation model is a cross-sectional regression. The unit of analysis is the province. The explanatory variable is the change in the share of immigrants over the period of analysis (2006-2015). The control variables include the change in (log) income per capita in female and male widows (%) over the period of analysis (2006-2015). Each regression is weighted by the province population 65+. The reported F Statistics is the Kleinbergen–Paap rk Wald F statistics. Robust standard errors in parenthesis. *p < 0.1 ** p < 0.05 *** p < 0.01


Next, we perform a falsification test to mitigate potential issues of reverse causality, examining whether the instrument-predicted change in the immigrant shares during our analysis period (2006-2015) explains changes in the main outcome variables in the pre-sample period (Table 10). We run a cross-sectional linear regression model, where the dependent variables are the changes in LR hospitalisations in column (1), Ac u t e cases in column (2), and the average LoS in column (3) in the period between 2001 and 2005. In column (4), relative to the mortality rate, the dependent variable refers to the period 2003-2005. The results indicate that the change in the instrument-predicted immigrant share in the 2006-2015 period, is not significantly correlated with changes in the outcomes considered in the pre-estimation period, suggesting that reverse causality is unlikely to constitute a threat to our identification strategy.


Table 10: Reverse Causality

	(1) Δ LR	(2) Δ Acute	(3) Δ LoS	(4) A Mortality
Variation of Predicted Immigration Share (2006-2015)	-0.000695	0.0737	6.597	125.6
	(0.0147)	(0.0869)	(7.065)	(162.1)
Observations	103	103	103	103
R-squared	0.048	0.072	0.083	0.060
Controls	YES	YES	YES	YES

Notes: The main explanatory variable is the variation of predicted immigration share between 2006 and 2015. The dependent variables are differentiated over the period 2001-2005 (Long-term and rehabilitation in (1) and acute cases in (2), the average length of stay for total hospitalisations in (3)) or 2003-2005 (total mortality for over 75 in (4)). The control variables at the province level include the (log) income per capita, female and male widows (%). Robust standard errors in parenthesis. * p < 0.05 *** p < 0.01

In Figure 4, we provide visual evidence to further examine the potential for reverse causality. The horizontal axis in each panel of the figure displays the change in the instrument-predicted immigrant share over the analysis period, while the vertical axis represents the variation in a preceding period for LR hospitalisations (2001-2005) in Panel a, average LoS of hospitalisations (2001-2005) in Panel b, and mortality (2003-2005) in Panel c. It is easy to see that there is no correlation between the pairs of variables, reinforcing our confidence in the absence of pre-trends among them.





c. Mortality

Notes: The x-axis reports the variation of predicted immigration share in the period 2006-2015. The y-axes report the variation in a pre-period for four main variables: the change in Long-term and rehabilitation hospitalisations between 2001 and 2005 (Figure a); the change in the average length of stay between 2001 and 2005 (Figure b); and the mortality rate for the population 75+ between 2003 and 2005 (Figure c).

One main concern when using the shift-share instrument is the potential endogeneity of the initial shares of the country-of-origin groups. To address this concern, we first assess the relevance of each country-of-origin share in generating the identifying variation in the instruments by calculating the Rotemberg weights, as in Goldsmith- Pinkham et al. (2020). The results are reported in Table 11. Panel A indicates that the weights are all positive confirming that all the shares are positively correlated with the instrument. Consequently, the estimated coefficient represents a convex combination of the individual country estimates, denoted as β_k . Panel B displays the correlation among the components of the IV estimates (g_k and z_k), the Rotemberg weights (α_k), the just-identified coefficient estimates (β_k), the first-stage F-statistic of the immigrant share (F_k), and the variation in the immigrant shares across locations (Var(z_k)). The correlation highlights the significance of both

the shares and the shifts in identifying variation. Panel C shows that immigrants from East Europe carry the largest weight. Lastly, Panel D reports statistics on how the values of β_k vary with the positive and negative Rotemberg weights.

Finally, in line with Goldsmith-Pinkham et al. (2020), we also test for any correlation between the initial shares of an immigrant group and the pre-period change in the outcome variables. The results presented in Table 12 indicate no significant correlation between the initial share for East Europe and the pre-period change.

Panel A: Negativ	e and positive we	ights			
-	Sum	Mea	an	Sh	nare
Negative	-0.000	-0.0	00	0.	000
Positive	1.000	0.10	67	1.	000
Panel B: Correlat	tions of Origin Ag	ggregates			
	α_k	g _k	β_k	F_k	$Var(z_k)$
α_k	1				
g _k	0.982	1			
β_k	-0.228	-0.228	1		
F _k	-0.004	-0.157	0.180	1	
$Var(z_k)$	0.855	0.777	-0.141	0.457	1
Panel C: Top 5 R	otemberg Weight	Origins			
	$\widehat{lpha}_{ m k}$	g_k	$\hat{\beta}_{\mathbf{k}}$	95 % CI	
East Europe	0.576	1.27e+06	-0.145	(-0.500,	0.010)
Asia	0.310	4.72e+05	-0.041	(-0.185,	0.135)
North Africa	0.027	1.45e+05	0.038	(-0.150,	0.500)
South America	0.050	97634.000	-0.045	(-0.190,	0.120)
Other Africa	0.037	1.36e+05	-0.059	(-0.195,	0.215)
Panel D: Estimate	es of β_k for Positi	ve and Negative	Weights		
	α-weighted Sum	Share o	f overall β		Mean
Negative	0.000	-(0.001		-0.131
Positive	-0.100	1	.001		-0.061

Table 11: Rotemberg weights

Notes: The table reports statistics on the Rotemberg weights (Goldsmith- Pinkham et al. 2020). Panel A reports the share and sum of negative weights. Panel B reports correlations between the weights (αk), the immigrant share (gk), the just-identified coefficient estimates (βk), the first-stage F-statistic of the immigrant share (Fk), and the variation in the immigrant shares across locations (Var(zk)). Panel C reports the top five nationalities according to the Rotemberg weights. The 95 percent confidence interval is the weak instrument robust confidence interval using the method from Chernozhukov & Hansen (2008) over a range from -10 to 10. Panel D reports statistics about how the values of β_k vary with the positive and negative Rotemberg weights.

	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
	Asia	East Europe	Other	North	Other	North	South	Oceania
Panel a	11010	Lust Lurope	Europe	Africa	Africa	America	America	orraina
Change in LR cases (2001-2005)	1.457	2.044	0.035	0.258	1.782	1.350	1.627	1.464
	(1.451)	(2.319)	(0.937)	(0.533)	(1.114)	(0.942)	(1.192)	(1.156)
Observations	91	91	91	91	91	91	91	91
R-squared	0.745	0.539	0.644	0.814	0.680	0.658	0.732	0.628
Controls	YES	YES	YES	YES	YES	YES	YES	YES
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
			Other	North	Other	North	South	
Panel b	Asia	East Europe	Europe	Africa	Africa	America	America	Oceania
Change in Acute cases								
(2001-2005)	0.111	0.689	0.019	-0.07	0.202	0.113	0.191	0.133
	-0.305	-0.503	-0.12	-0.093	-0.215	-0.183	-0.253	-0.209
Observations	91	91	91	91	91	91	91	91
R-squared	0.742	0.56	0.644	0.815	0.674	0.652	0.73	0.622
Controls	YES	YES	YES	YES	YES	YES	YES	YES
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
			Other	North	Other	North	South	
Panel c	Asia	East Europe	Europe	Africa	Africa	America	America	Oceania
Change in avg LoS (2001-2005)	0.002	0.013	0.000	0.004*	0.006	0.002	0.004	0.005
	(0.009)	(0.011)	(0.003)	(0.002)	(0.006)	(0.005)	(0.007)	(0.006)
Observations	91	91	91	91	91	91	91	91
R-squared	0.742	0.561	0.644	0.823	0.682	0.653	0.731	0.629
Controls	YES	YES	YES	YES	YES	YES	YES	YES

Table 12: Exogeneity of the Initial (1991) Shares of Country-of-Origin Groups, OLS

	(1)	(2)	(3) Other	(4) North	(5) Other	(6) North	(7) South	(8)
Panel d	Asia	East Europe	Europe	Africa	Africa	America	America	Oceania
Change in mortality (2003-2005)	-0.000 (0.001)	0.001 (0.001)	0.000 (0.001)	-0.000 (0.000)	-0.001 (0.001)	0.000 (0.001)	-0.000 (0.001)	0.001 (0.001)
Observations	91	91	91	91	91	91	91	91
R-squared	0.741	0.533	0.644	0.814	0.669	0.649	0.726	0.621
Controls	YES	YES	YES	YES	YES	YES	YES	YES

Notes: The estimation model is a cross-sectional regression. The unit of analysis is the province. The dependent variables are the 1991 immigrant shares by area of origin. The explanatory variable is the change in Long-term and rehabilitation hospitalisations over the years 2001-2005 in panel a; the change in acute hospitalisations over the years 2001-2005 in panel b, the change in the average length of stay in panel c and the change in mortality (for the population 75+) over the years 2003-2005 in panel d. The controls at the province level include income pc (log), female and male widows (%). All the controls are the 2006 value. Each regression is weighted by the over-65 Italian population of the province. Robust standard errors in parenthesis with * p < 0.10, ** p < 0.05, *** p < 0.01.

28

5.3.2 Pre-Trend Analysis

We further ensure the robustness of our estimates by accounting for existing pre-trends in hospitalisations and average length of stay, following the approach of Dix-Carneiro et al. (2018). As depicted in Table 13, we enrich our model specification by including the change in the dependent variable in the pre-sample period as an additional control variable. Consistently with the estimates in Table 4, Table 5, and Table 6, the impact of immigration on Long-term and rehabilitation hospitalisations (in column 1), the average length of stay (column 3), and total mortality (column 4) remains negative and significant. Conversely, no effect is detected for acute cases (column 2).

	(1)	(2)	(3)	(4)
	LR	AC	LoS	Mortality
Share of immigrants	-0.124***	0.208	-8.093***	-183.9*
e	(0.0241)	(0.218)	(2.953)	(107.8)
Δ LR	-0.0875**	(0.2.0)	()	()
	(0.0364)			
Δ Acute	. /	-0.0200		
		(0.0286)		
Δ LoS			-0.0398***	
			(0.00688)	
Δ Mortality				0.01000
				(0.00907)
Observations	1,030	1,030	1,030	1,030
R-squared	0.129	0.686	0.293	0.440
Number of provinces	103	103	103	103
Controls	YES	YES	YES	YES
Year FE	YES	YES	YES	YES
Province FE	YES	YES	YES	YES
F Stat	36.688	45.650	22.606	38.481

Table 13: Pre-Trend Analysis (Dix-Carneiro et al., 2018)

Notes: The dependent variables are: Long-term and rehabilitation hospitalisations (column 1), acute cases (column 2), average length of stay (column 3) and mortality for population 75+ (column 4). The main explanatory variable is the ratio between the working-age immigrants over the native population 65+. The controls at the province level include income pc (log), female and male widows (%). All the controls are the 2006 value. In these specifications, we added to the above-mentioned control variables the variation in the outcomes considered for the period 2001-2005 (the only exception is mortality, for which we used the period 2003-2005 due to data limitations). These variations are interacted with a trend variable. Each specification includes

year and province fixed effect. The reported F Statistics is the Kleinbergen–Paap rk Wald F statistics. Robust standard errors in parenthesis with p < 0.1 * p < 0.05 * p < 0.01

5.4 Home-based care and public health spending

In this section, we conduct a preliminary assessment of the potential reduction in public health expenditures arising from an increase in the supply of non-specialized home-based care provided by immigrants. Given that the IV coefficient estimates may not necessarily represent average treatment effects for the entire Italian population, these calculations are intended as a back-of-the-envelope exercise designed to exemplify the implications of the estimates for public healthcare spending.

Our main findings, presented in Table 4, suggest that a rise in the ratio of working-age migrants to the population aged 65 and over by 0.116 (from 0.169 to 0.285) over the analysis period, equivalent to 0.013 annually, generated a reduction of long-term and rehabilitation hospitalisations of 17,948 cases, representing a 5% decrease relative to the baseline.¹²

Given these findings, we assess the potential cost savings resulting from reduced hospitalisations attributed to migrant-provided home-based care. To quantify the costs associated with hospitalisations, we use Diagnosis Related Group (DRG) rates, a system that categorises hospital admissions by allocating specific costs and standard durations of hospitalisation to each admission. While the DRG rates are established nationally, regional variations exist due to healthcare decentralisation, allowing regions to adjust rates based on local strategies, cost of living, or infrastructure needs. As our analysis primarily focuses on quantifying the monetary consequences of home-based care provision at the national level, we rely on the national DRG rates. In particular, we consider hospitalisation costs for each illness group and hospitalisation type, focusing on long-term and rehabilitation admissions.

In our context, supported by the evidence presented in the paper, savings in secondary healthcare expenditure result from two distinct channels. Firstly, migrant-provided home-based care reduces the occurrence of long-term hospitalisations, impacting the extensive margin. Secondly, our findings demonstrate that, in addition to the reduced hospitalisation frequency, migrant home-based care also leads to shorter average hospital stays, influencing the intensive margin. To provide an accurate assessment of the total reduction, both effects must be taken into account.

¹² The reduction of 17,948 is computed by first multiplying the relevant coefficient (-0.118 in Table 4) by 0.013, and then by the average annual size of the Italian native elderly population, i.e. 11,700,000. The 5% share is computed by dividing the variation in the number of cases over the (average) total annual number of long-term and rehabilitation hospitalisations at the national level 327,932.

Table 14: Back-of-the-envelope calculations of potential savings due to a reduction in Long-termcare and rehabilitation hospitalisations

Disorder	Prevalence	Cost	Coef.	Estimated	Estimated	Estimated	Estimated
group		per	estimate	savings	savings	savings	savings
		hospitalisation	(# of cases)	(100k)	2007	2023	2050
					(11.7 mln)	(14.2 mln)	(18.9 mln)
Genitourinary	0.00727	7762	-0.051000	-514,588	-60,206,786	-73,071,484	-97,338,421
Disord.							
Circulatory	0.00588	9251					
Disord.							
Traumatic	0.00296	9312	-0.019200	-232,416	-27,192,630	-33,003,021	-43,963,277
Injuries							
Musculoskelet	0.00303	6371	-0.032600	-270,018	-31,592,124	-38,342,578	-51,076,094
Disord.							
Respiratory	0.00084	5648					
Disord.							
Nervous Sys.	0.00174	9700	-0.015000	-189,154	-22,131,036	-26,859,890	-35,780,021
Disord.							
Cancers	0.00069	7043					
Mental Disord.	0.00043	5575	-0.003160	-22,903	-2,679,677	-3,252,257	-4,332,328
Total				-1,229,079	-143,802,252	-174,529,229	-232,490,141

Notes: The costs are expressed in current euro and are defined according to reimbursement rates registered in 2011. Population estimates refers to the 65+ Italian residents sourced from the ISTAT statistics and population projections. We assume a rise in the ratio of working-age migrants to the population aged 65 and over 1.3 percentage points annually, resulting from the overall rise by 11.6 percentage points over the analysis period.

The results of the back-of-the-envelope calculations on potential savings are presented in Table 14 and Table 15. As mentioned, these calculations refer to two channels, namely a reduced number and a reduced duration of hospitalisations. We specifically address LR stays, where our analysis identified significant effects of immigrant home-based care provision. We focus on the eight most frequent hospitalisation types, as detailed in the second column of Table 14. Notably, nervous system disorders, circulatory disorders, and traumatic injuries exhibit the highest costs (euros 9700, 9251, and 9312, respectively), as indicated in the third column of the Table.

Utilising the coefficient estimates indicating the reduction in the number of hospitalisations for each diagnostic group, as presented in Table 7 and detailed in the fourth column of Table 14, we calculate the potential savings on the extensive margin. This calculation assumes a 1.3 percentage point annual increase in the ratio between the number of working-age immigrants and the native population aged 65 and older, as evidenced in the analysis period considered. With the coefficient estimates and unit costs of admissions considered, the most substantial savings per 100 thousand residents are observed

in genitourinary, musculoskeletal, and trauma-related hospitalisations, amounting to 515, 270, and 232 thousand euros annually, respectively. The cumulative savings across all categories would total 1,229 thousand euros annually.

To contextualise this figure, we scale these potential savings by the elderly population size, which stood at 11.7 million residents aged 65 and older in 2007, resulting in an annual savings of 143.8 million euros (sixth column in Table 14). Given the rapid aging of the Italian population, these savings may increase to 174.5 million euros based on current population estimates (14.2 million in 2023) and further to 232.5 million euros in 2050, based on ISTAT population projections, foreseen to rise to 18.9 million 65+ individuals by 2050.

Disorder	Prevalence	Cost	Coef.	Estimated	Estimated	Estimated	Estimated
group		per	estimate	savings	savings	savings	savings
		hospitalisation	(# of cases)	(100k)	2007	2023	2050
					(11.7 mln)	(14.2 mln)	(18.9 mln)
Genitourinary	28.41	381	-91.83	-330,748	-38,697,534	-46,966,237	-62,563,659
Disord.							
Circulatory	28.40	499	-53.98	-205,554	-24,049,866	-29,188,726	-38,882,261
Disord.							
Traumatic	31.08	378	-106.00	-154,477	-18,073,864	-21,935,801	-29,220,650
Injuries							
Musculoskelet	24.50	379	-59.43	-88,768	-10,385,887	-12,605,093	-16,791,227
Disord.							
Respiratory	25.01	339	-120.20	-44,555	-5,212,944	-6,326,821	-8,427,949
Disord.	10.00						
Nervous Sys.	40.00	351	-47.87	-37,945	-4,439,604	-5,388,237	-7,177,663
Disord.	24.46	457					
Cancers	24.46	457		-	-	-	-
Mental Disord.	33.62	189		-	-	-	-
Total				-862,049	-100,859,698	-122,410,916	-163,063,409

Table 15: Back-of-the-envelope calculations of potential savings due to a reduction in the lengthof Long-term care and rehabilitation hospitalisations

Notes: The costs are expressed in current euro and are defined according to reimbursement rates registered in 2011. Population estimates refers to the 65+ Italian residents sourced from the ISTAT statistics and population projections. We assume a rise in the ratio of working-age migrants to the population aged 65 and over 1.3 percentage points annually, resulting from the overall rise by 11.6 percentage points over the analysis period.

Beyond the extensive margin considerations, it is crucial to incorporate potential savings stemming from a decrease in the duration of long-term and rehabilitation stays (intensive margin). While certain hospitalisations may be entirely avoided for individuals with home-based care, others, though occurring, are likely to have shorter durations as care can be administered in a home setting.

In Table 15, we present the daily costs of hospitalisations (third column) and the corresponding coefficient estimates presented in Table 8, reiterated here in the fourth column. Given these coefficient estimates and assuming a one percentage point increase in the ratio between the number of working-age immigrants and the native population aged 65+, coupled with the prevalence of respective hospitalisations outlined in the second column of Table 14, the potential savings for 100 thousand elderly individuals would predominantly arise from genitourinary hospitalisations, and overall total saving would culminate in an annual reduction of 862 thousand euros.

Considering the size of the elderly population at various time points (2007, 2023, and 2050), the corresponding savings at the intensive margin would be 100.9 million, 122.4 million, and 163.1 million euros, respectively.

Taking into account both the intensive and extensive margins, the potential secondary care savings in 2007 could amount to 244.6 million euros. This constitutes approximately 8% of the SDO costs allocated to LR elderly hospitalisations for the eight diagnostic groups considered, totaling 3,122.5 million euros in 2007. In 2023, the potential savings in secondary care could amount to 292.9 million euros. Considering that total public spending on hospitalisations amounted to 50,135 million euros in 2022, these savings represent approximately 0.59% of the total relevant annual public expenditure.¹³ Looking at the long-term perspective, especially considering Italy's significant pace of population ageing, these potential savings are likely to become increasingly important and could rise by a third.

6 Conclusions

The arrival of immigrants specialising in the provision of domestic services is likely to reshape the time allocation of households, influencing a wide array of outcomes ranging from the business cycle to fertility decisions (Mariani and Rosati, 2022). Yet, one aspect overlooked in the literature is the impact on the welfare of the elderly resulting from the increased availability of paid non-specialized home-based care. With a growing share of the elderly in the population and the weakening of the family networks, both institutionalised and informal elderly care are put under pressure. By complementing or substituting for formal care and/or informal care provided by relatives, the home-based care provided by immigrants could potentially improve the welfare of the elderly and may also yield benefits for the public finances.

In this paper, we provide novel causal evidence on the effects of immigration on elderly health, focusing on hospitalisations, their duration, and mortality rates. We employ an instrumental variable

¹³ This data is taken from the system of health account SHA, by ISTAT.

approach to tackle the potential endogeneity of immigrants' location and extensively validate our instrument choice. Our results show that the arrival of immigrants specialising in domestic services reduces hospitalisation rates and their duration, particularly for long-term cases. Moreover, these effects extend to mortality rates, particularly among females, indicating that the presence of immigrants not only reduces the use of formal care, but also contributes to overall improvements in elderly health.

Disaggregating our analysis by diagnosis types, we find that our baseline results in terms of longterm and rehabilitation hospitalisation frequency are driven by mental, nervous system, genitourinary, musculoskeletal, and traumatic injury cases. Additionally, concerning the average length of stay, we identify widespread reductions across several disease categories, up to 1.2 days for disorders related to the respiratory system.

Lastly, we assess the potential cost savings resulting from the decline in LR hospitalisations due to the presence of immigrant home-based care. Using our estimates of the decrease in the frequency of hospitalisations for each diagnostic group, we quantify potential savings on the extensive margin. Subsequently, we incorporate potential savings on the intensive margin, stemming from the accompanying reduction in the length of stay associated with these hospitalisations. These findings underscore the significance of home-based care during the post-discharge phase, particularly highlighting the valuable contribution of immigrant caregivers in assisting the elderly. Simultaneously, they point to substantial reductions in public health spending on formal care. A rough estimation, conducted through a back-of-the-envelope calculation, suggests that the registered annual increase in the migrant-to-elderly population ratio (averaging 1.3 percentage points annually during our analysis period) corresponds to roughly 8% of the LR hospitalisations dedicated to the elderly observed in the study. Extrapolating from this, it's plausible to infer that the arrival of immigrants over our study period may have yielded annual savings to the public budget, amounting to approximately 0.59% of total expenditures on hospitalisation or about 9 per cent over our study period.

References

- Altonji, J. G., & Card, D. (1991). The Effects of Immigration on the Labor Market Outcomes of Less-Skilled Natives. In *Immigration, Trade and the Labor Market* (pp. 201–234). NBER.
- Barnay, T., & Juin, S. (2016). Does home care for dependent elderly people improve their mental health? *Journal of Health Economics*, 45, 149–160. https://doi.org/10.1016/j.jhealeco.2015.10.008
- Bettin, G., & Sacchi, A. (2020). Health spending in Italy: The impact of immigrants. *European Journal* of Political Economy, 65. https://doi.org/10.1016/j.ejpoleco.2020.101932
- Borusyak, K., Hull, P., & Jaravel, X. (2022). Quasi-Experimental Shift-Share Research Designs. *Review* of Economic Studies, 89(1), 181–213. https://doi.org/10.1093/restud/rdab030
- Byrne, D., Goeree, M. S., Hiedemann, B., & Stern, S. (2009). Formal Home Health Care, Informal Care, and Family Decision Making. *International Economic Review*, *50*(4).
- Card, D. (2001). Immigrant inflows, native outflows, and the local labor market impacts of higher immigration. *Journal of Labor Economics*, 19(1), 22–64. https://doi.org/10.1086/209979
- Charles, K. K., & Sevak, P. (2005). Can family caregiving substitute for nursing home care? *Journal of Health Economics*, 24(6), 1174–1190. https://doi.org/10.1016/j.jhealeco.2005.05.001
- Chernozhukov, V., & Hansen, C. (2008). The reduced form: A simple approach to inference with weak instruments. *Economics Letters*, 100(1), 68–71. https://doi.org/10.1016/j.econlet.2007.11.012
- Costa-Font, J., Jimenez-Martin, S., & Vilaplana, C. (2018). Does long-term care subsidization reduce hospital admissions and utilization? *Journal of Health Economics*, 58, 43–66. https://doi.org/10.1016/j.jhealeco.2018.01.002
- Dix-Carneiro, R., Soares, R. R., & Ulyssea, G. (2018). Economic shocks and crime: Evidence from the Brazilian trade liberalization. *American Economic Journal: Applied Economics*, 10(4), 158–195. https://doi.org/10.1257/app.20170080
- DOMINA. (2022). 4° Rapporto Annuale sul Lavoro Domestico. www.domesticworkobservatory.com
- Furtado, D., & Ortega, F. (2023). Does Immigration Improve Quality of Care in Nursing Homes? *Journal* of Human Resources.
- Goldsmith-Pinkham, P., Sorkin, I., & Swift, H. (2020). Bartik instruments: What, when, why, and how. *American Economic Review*, *110*(8), 2586–2624. https://doi.org/10.1257/AER.20181047
- Grabowski, D. C., Gruber, J., & Mcgarry, B. (2023). *Immigration, The Long-Term Care Workforce, and Elder Outcomes in the U.S.*
- Grossman, M. (1972). On the Concept of Health Capital and the Demand for Health. *The Journal of Political Economy*, 80(2), 223–255.
- Kemper, P., Weaver, F., Short, P. F., Shea, D., & Kang, H. (2008). Meeting the need for personal care among the elderly: Does medicaid home care spending matter? *Health Services Research*, 43(1 P2), 344–362. https://doi.org/10.1111/j.1475-6773.2007.00762.x
- Lean, K., Nawaz, R. F., Jawad, S., & Vincent, C. (2019). Reducing urinary tract infections in care homes by improving hydration. *BMJ Open Quality*, 8(3). https://doi.org/10.1136/bmjoq-2018-000563
- Mariani, R. D., Pasquini, A., & Rosati, F. C. (2023). The Immigration Puzzle in Italy: A Survey of Evidence and Facts. In *Italian Economic Journal* (Vol. 9, Issue 1, pp. 85–116). Springer Science and Business Media Deutschland GmbH. https://doi.org/10.1007/s40797-021-00168-x

- Mariani, R. D., & Rosati, F. C. (2022). Immigrant supply of marketable child care and native fertility in Italy. *Journal of Demographic Economics*, 88(4), 503–533. https://doi.org/10.1017/dem.2021.28
- Melchiorre, M. G., Quattrini, S., Lamura, G., & Socci, M. (2021). A mixed-methods analysis of care arrangements of older people with limited physical abilities living alone in Italy. *International Journal of Environmental Research and Public Health*, 18(24). https://doi.org/10.3390/ijerph182412996
- NNA. (2021). L'assistenza agli anziani non autosufficienti in Italia 7° Rapporto. www.maggiolieditore.it
- Peri, G., Romiti, A., & Rossi, M. (2015). Immigrants, domestic labor and women's retirement decisions. *Labour Economics*, *36*, 18–34. https://doi.org/10.1016/j.labeco.2015.07.004
- Rice, J. B., Kasper, J. D., & Pezzin, L. E. (2009). A comparative analysis of medicaid long-term care policies and their effects on elderly dual enrollees. *Health Economics*, 18(3), 275–290. https://doi.org/10.1002/hec.1367
- Rogers, M. A. M., Fries, B. E., Kaufman, S. R., Mody, L., McMahon, L. F., & Saint, S. (2008). Mobility and other predictors of hospitalization for urinary tract infection: A retrospective cohort study. *BMC Geriatrics*, 8. https://doi.org/10.1186/1471-2318-8-31
- Van Houtven, C. H., & Norton, E. C. (2004). Informal care and health care use of older adults. *Journal of Health Economics*, 23(6), 1159–1180. https://doi.org/10.1016/j.jhealeco.2004.04.008
- Weaver, F. M., & Weaver, B. A. (2014). Does availability of informal care within the household impact hospitalisation? *Health Economics, Policy and Law, 9*(1), 71–93. https://doi.org/10.1017/S1744133113000169

A Appendices

A.1 Descriptive statistics by diagnostic group

Table A1 shows the descriptive statistics for long-term and rehabilitation cases by the most prevalent major diagnostic groups considered in the analysis.

Variable	Obs.	Mean	St. Dev.
Panel a: Hospitalisa	itions		
Mental	1030	.000428	.0004453
Cancers	1030	.000693	.0009226
Nervous	1030	.001736	.0012235
Circulatory	1030	.005875	.0032728
Respiratory	1030	.000841	.0008245
Genitourinary	1030	.007266	.0040977
Musculoskeletal	1030	.003033	.0021787
Traumatic	1030	.002962	.0021208
Panel b: Average Lo	<i>S</i>		
Mental	1030	33.619	17.045
Cancers	1030	24.461	9.702
Nervous	1030	40.000	8.726
Circulatory	1030	28.399	6.786
Respiratory	1030	25.013	5.766
Genitourinary	1030	28.412	6.786
Musculoskeletal	1030	24.500	6.829
Traumas	1030	31.076	7.318

Table A1: Descriptive statistics for LR cases by disease

Notes: The data refer to 103 provinces spanning the time period from 2006 to 2015. Statistics are weighted by the relevant province 65+ population size.

A.2 OLS results

Table A2.1, Table A2.2 and Table A2.3 report the results of the OLS regression.

	(1)	(2)	(3)
	Total	Acute	LR
Share of immigrants	0.219	0.237*	-0.0181
C	(0.140)	(0.133)	(0.0210)
Income pc(log)	0.0635	0.0515	0.0119
	(0.128)	(0.116)	(0.0126)
Widowers (percent)	0.0652***	0.0596***	0.00559
	(0.0216)	(0.0211)	(0.00388)
Widows (percent)	0.0114	0.0135	-0.00210
- · ·	(0.0286)	(0.0303)	(0.00337)
Observations	1,030	1,030	1,030
R-squared	0.618	0.634	0.042
Number of provinces	103	103	103
Year FE	YES	YES	YES
Province FE	YES	YES	YES
	(1 1 1)	1	1 (()) 771

Table A2.1: OLS results for hospitalisation

Notes: OLS Estimates (including year and province fixed effects). The dependent variables are: total hospitalisations over native population 65+ (column 1), acute hospitalisations over native population 65+ (column 2), long-term and rehabilitation hospitalisations over native population 65+ (column 3). The explanatory variable is the ratio between the working age immigrants over the native population 65+. Robust standard errors in parenthesis with * p<0.1 ** p<0.05 *** p<0.01

	(1)	(2)	(3)
	Total	Acute	LR
Share of immigrants	-4.407**	-2.343*	-16.15
C	(1.929)	(1.378)	(11.78)
Income pc(log)	-0.422	-0.555	-2.167
	(0.631)	(0.465)	(2.976)
Widowers (percent)	-0.672*	-0.739***	-3.822
-	(0.378)	(0.257)	(2.628)
Widows (percent)	-0.532	-0.273	-0.430
	(0.611)	(0.414)	(1.352)
Observations	1,030	1,030	1,030
R-squared	0.157	0.181	0.065
Number of provinces	103	103	103
Year FE	YES	YES	YES
Province FE	YES	YES	YES

Table A2.2: OLS results for average LoS

Notes: OLS Estimates (including year and province fixed effects). The dependent variables are: average LoS for total hospitalisations (column 1), for acute hospitalisations (column 2), for long-term and rehabilitation hospitalisations (column 3). The explanatory variable is the ratio between the working age immigrants over the native population 65+. Robust standard errors in parenthesis with p < 0.1 * p < 0.05 * p < 0.01

	(1)	(2)	(3)
	Mortality(tot)	Mortality(f)	Mortality(m)
Share of immigrants	-49.49	-69.84	29.44
	(72.97)	(65.47)	(75.48)
Income pc(log)	-43.70	-63.51	-31.24
	(45.51)	(40.25)	(54.88)
Widowers (percent)	-12.80	-6.437	-21.25
-	(15.88)	(15.02)	(19.97)
Widows (percent)	-22.67	-17.22	-27.14*
	(15.87)	(12.94)	(14.76)
Observations	1,030	1,030	1,030
R-squared	0.333	0.402	0.294
Number of provinces	103	103	103
Year FE	YES	YES	YES
Province FE	YES	YES	YES

Table A2.3: OLS results for mortality

Notes: OLS Estimates (including year and province fixed effects). The dependent variables are: total mortality for the population 75+ (column 1), mortality for the female population 75+ (column 2), mortality for the male population 75+ (column 3). The explanatory variable is the ratio between the working age immigrants over the native population 65+. Robust standard errors in parenthesis with p < 0.1 * p < 0.05 * p < 0.01

A.3 Adding a control for informal care provided by family members

In order to control for the fact that some households may choose to use one of the household members as informal caregiver, we add the share of women aged 45-64 over the Italian population 65+ in the main regression. As the table below shows, the results do not change.

	(1)	(6)	(7)
	(1)	(6)	(7)
	Acute and	Acute	LR
	LR		
Share of immigrants	0.0801	0.187	-0.107***
Share of miningrants		01201	
	(0.232)	(0.228)	(0.0247)
Observations	1,030	1,030	1,030
R-squared	0.675	0.685	0.147
Number of provinces	103	103	103
Controls	YES	YES	YES
Year FE	YES	YES	YES
Province FE	YES	YES	YES
F Stat	36.669	36.669	36.669

Table A3.1: Hospitalisations (with share of women 45-64 years old as additional control)

Notes: Estimates of Equation 2 (including year and province fixed effects) with the addition of the share of women 45-64 years as control variable. In column 1 we include all hospitalisations, in column 2 we focus on acute hospitalisations, and in column 3 on long-term and rehabilitation. The endogenous regressor is the ratio between the working-age immigrant resident population over the size of native population aged 65 and older. Each regression is weighted by the province population 65+. Each regression is weighted by the province population 65+. The reported F Statistics is the Kleinbergen–Paap rk Wald F statistics. Robust standard errors in parenthesis with * p < 0.1 ** p < 0.05 *** p < 0.01

	(1)	(6)	(7)
	Acute and	Acute	LR
	LR		
Share of immigrants	-13.67***	-2.487*	-102.9***
-	(2.269)	(1.304)	(20.15)
Observations	1,030	1,030	1,030
	0.094	0.162	-0.068
R-squared			
Number of provinces	103	103	103
Controls	YES	YES	YES
Year FE	YES	YES	YES
Province FE	YES	YES	YES
F Stat	36.669	36.669	36.669

Table A3.2: Average LoS (with share of women 45-64 years old as additional control)

Notes: Estimates of Equation 2 (including year and province fixed effects) with the addition of the share of women 45-64 years as control variable. In column 1 we refer to all hospitalisations, in column 2 to acute hospitalisations, while in column 3 to long-term and rehabilitation hospitalisations. The endogenous regressor is the ratio between the working-age immigrant resident population over the size of native population aged 65 and older. Each regression is weighted by the province population 65+. The reported F Statistics is the Kleinbergen–Paap rk Wald F statistics. Robust standard errors in parenthesis with * p<0.1** p<0.05*** p<0.01

Table A3.3: Mortality (with share of women 45-64 years old as additional control)

	(1)	(6)	(7)
	Total	Female	Male
Share of immigrants	-117.9	-173.8	-61.96
	(106.3)	(112.3)	(115.7)
Observations	1,030	1,030	1,030
R-squared	0.465	0.474	0.445
Number of provinces	103	103	103
Controls	YES	YES	YES
Year FE	YES	YES	YES
Province FE	YES	YES	YES
F Stat	36.669	36.669	36.669

Notes: Estimates of Equation 2 (including year and province fixed effects) with the addition of the share of women 45-64 years as control variable. In column 1 we refer to standardized mortality rate of total population 75+, in column 2 to female population 75+, while in column 3 to male population 75+. The endogenous regressor is the ratio between the working-age immigrant resident population over the size of native population aged 65 and older. Each regression is weighted by the province population 65+. The reported F Statistics is the Kleinbergen–Paap rk Wald F statistics. Robust standard errors in parenthesis with * p<0.1 * * p<0.05 *** p<0.01

RECENT PUBLICATIONS BY CEIS Tor Vergata

The Cost of Coming Out

Enzo Brox and Riccardo Di Francesco CEIS *Research Paper*, 572 April 2024

The Time-Varying Multivariate Autoregressive Index Model Gianluca Cubadda, Stefano Grassi and Barbara Guardabascio CEIS *Research Paper*, 571 January 2024

A Further Look at the Gender Gap in Italian Academic Careers Marianna Brunetti, Annalisa Fabretti and Mariangela Zoli CEIS *Research Paper*, 570 December 2023

The Wind of Populism: Voter Turnout and Political Distance Leonardo Becchetti and Gianluigi Conzo CEIS *Research Paper*, 569 December 2023

Funding Liquidity and Stocks' Market Liquidity: Structural Estimation From High-Frequency Data

Gian Piero Aielli and Davide Pirino CEIS *Research Paper*, 568 November 2023

On The Nonlinearity of the Finance and Growth Relation: the Role of Human Capital Alberto Bucci, Boubacar Diallo and Simone Marsiglio CEIS *Research Paper*, 567 November 2023

Distance Work and Life Satisfaction after the COVID-19 Pandemics Leonardo Becchetti, Gianluigi Conzo and Fabio Pisani CEIS *Research Paper*, 566 September 2023

Scitovsky Was Right...and There Is More: Comfort Goods, Stimulus Goods, Education and Subjective Wellbeing

Leonardo Becchetti and Chiara Lubicz CEIS Research Paper, 565 August 2023

Information Campaigns and migration perceptions

Erminia Florio *CEIS Research Paper, 564* August 2023

Is Self-Employment for Migrants? Evidence from Italy

Marianna Brunetti and Anzelika Zaiceva *CEIS Research Paper, 563* July 2023

DISTRIBUTION

Our publications are available online at <u>www.ceistorvergata.it</u>

DISCLAIMER

The opinions expressed in these publications are the authors' alone and therefore do not necessarily reflect the opinions of the supporters, staff, or boards of CEIS Tor Vergata.

COPYRIGHT

Copyright © 2024 by authors. All rights reserved. No part of this publication may be reproduced in any manner whatsoever without written permission except in the case of brief passages quoted in critical articles and reviews.

MEDIA INQUIRIES AND INFORMATION

For media inquiries, please contact Barbara Piazzi at +39 06 72595652/01 or by email at <u>piazzi@ceis.uniroma2.it</u>. Our web site, www.ceistorvergata.it, contains more information about Center's events, publications, and staff.

DEVELOPMENT AND SUPPORT

For information about contributing to CEIS Tor Vergata, please contact at +39 06 72595601 or by e-mail at <u>segr.ceis@economia.uniroma2.it</u>